

Dejean's conjecture holds for $n \geq 30$

James Currie* and Narad Rampersad†
 Department of Mathematics and Statistics
 University of Winnipeg
 515 Portage Avenue
 Winnipeg, Manitoba R3B 2E9 (Canada)
 j.currie@uwinnipeg.ca
 n.rampersad@uwinnipeg.ca

January 6, 2009

Abstract

We extend Carpi's results by showing that Dejean's conjecture holds for $n \geq 30$.

1 Introduction

Repetitions in words have been studied starting with Thue [12, 13] at the beginning of the previous century. Much study has also been given to repetitions with fractional exponent [1, 3, 4, 5, 6, 8]. If $n > 1$ is an integer, then an **n -power** is a non-empty word x^n , i.e., word x repeated n times in a row. For rational $r > 1$, a **fractional r -power** is a non-empty word $w = x^{\lfloor r \rfloor} x'$ such that x' is the prefix of x of length $(r - \lfloor r \rfloor)|x|$. For example, 01010 is a $5/2$ -power. A basic problem is that of identifying the *repetitive threshold* for each alphabet size $n > 1$:

What is the infimum of r such that an infinite sequence on n letters exists, not containing any r -powers?

*The author is supported by an NSERC Discovery Grant.

†The author is supported by an NSERC Postdoctoral Fellowship.

We call this infimum the *repetitive threshold* of an n -letter alphabet, denoted by $RT(n)$. Dejean's conjecture [4] is that

$$RT(n) = \begin{cases} 7/4, & n = 3 \\ 7/5, & n = 4 \\ n/(n-1) & n \neq 3, 4 \end{cases}$$

The values $RT(2)$, $RT(3)$, $RT(4)$ were established by Thue, Dejean and Pansiot, respectively [12, 4, 11]. Moulin-Ollagnier [10] verified Dejean's conjecture for $5 \leq n \leq 11$, while Mohammad-Noori and Currie [9] proved the conjecture for $12 \leq n \leq 14$.

An exciting new development has recently occurred with the work of Carpi [3], who showed that Dejean's conjecture holds for $n \geq 33$. Verification of the conjecture is now only lacking for a finite number of values. In the present paper, we sharpen Carpi's methods to show that Dejean's conjecture holds for $n \geq 30$.

2 Preliminaries

The following definitions are from sections 8 and 9 of [3]: Fix $n \geq 30$. Let $m = \lfloor (n-3)/6 \rfloor$. Let $A_m = \{1, 2, \dots, m\}$. Let $\ker \psi = \{v \in A_m^* \mid \forall a \in A_m, 4 \text{ divides } |v|_a\}$. (In fact, this is not Carpi's *definition* of $\ker \psi$, but rather the assertion of his Lemma 9.1.) A word $v \in A_m^+$ is a **ψ -kernel repetition** if it has period q and a prefix v' of length q such that $v' \in \ker \psi$, $(n-1)(|v|+1) \geq nq-3$.

It will be convenient to have the following new definition: If v has period q and its prefix v' of length q is in $\ker \psi$, we say that q is a **kernel period** of v .

As Carpi states at the beginning of section 9 of [3]:

By the results of the previous sections, at least in the case $n \geq 30$, in order to construct an infinite word on n letters avoiding factors of any exponent larger than $n/(n-1)$, it is sufficient to find an infinite word on the alphabet A_m avoiding ψ -kernel repetitions.

For $m = 5$, Carpi produces such an infinite word, based on a paper-folding construction. He thus establishes Dejean's conjecture for $n \geq 33$. In the present paper, we give an infinite word on the alphabet A_4 avoiding ψ -kernel repetitions. We thus establish Dejean's conjecture for $n \geq 30$.

Definition 1. Let $f : A_4^* \rightarrow A_4^*$ be defined by $f(1) = 121$, $f(2) = 123$, $f(3) = 141$, $f(4) = 142$. Let w be the fixed point of f .

It is useful to note that the frequency matrix of f , i.e.,

$$[|f(i)|_j]_{4 \times 4} = \begin{bmatrix} 2 & 1 & 0 & 0 \\ 1 & 1 & 1 & 0 \\ 2 & 0 & 0 & 1 \\ 1 & 1 & 0 & 1 \end{bmatrix}$$

has an inverse modulo 4.

Remark 1. Let q be a non-negative integer, $q \leq 1966$. Fix $n = 32$.

R1: Word w contains no ψ -kernel repetition v with kernel period q .

R2: Word w contains no factor v with kernel period q such that $|v|/q \geq 35/34$.

Note that $\frac{32}{31} - \frac{34}{31q} = \frac{35}{34}$ when $q = \frac{34^2}{3} = 385\frac{1}{3}$, so neither piece of the remark implies the other. Note also that the conditions of the remark become **less** stringent for $n = 30, 31$. One also verifies that

$$\frac{35}{34} + \frac{9}{2(1967)} \leq \frac{32}{31} - \frac{34}{31q}$$

for $q \geq 1967$. To show that w contains no ψ -kernel repetitions for $n = 30, 31, 32$, it thus suffices to verify R1 and to show that word w contains no factor v with kernel period $q \geq 1967$ such that

$$|v|/q \geq 35/34 + 9/2(1967). \tag{1}$$

The remarks R1 and R2 are verified by computer search, so we will consider the second part of this attack. Fix $q \geq 1967$, and suppose that v is a factor of w with kernel period q , and $|v|/q \geq 35/34$. Without loss of generality, suppose that no extension of v has period q . Write $v = sf(u)p$ where s (resp. p) is a suffix (resp. prefix) of the image of a letter, and $|s|$ (resp. $|p|$) ≤ 2 .

If $|v| \leq q + 2$, then $35/34 \leq (q + 2)/q$ and $1/34 \leq 2/q$, forcing $q \leq 68$. This contradicts R2. We will therefore assume that $|v| \geq q + 3$.

Suppose $|s| = 2$. Since $|v| \geq q + 3$, write $v = s0zs0v'$, where $|s0z| = q$. Examining f , we see that the letter a_s preceding any occurrence of $s0$ in w is

uniquely determined if $|s| = 2$. It follows that $a_s v$ is a factor of w with kernel period q , contradicting the maximality of v . We conclude that $|s| \leq 1$.

Again considering f , we see that if t is any factor of w of length 3, and $u_1 t$, $u_2 t$ are prefixes of w , then $|u_1| \equiv |u_2| \pmod{3}$. Since $|v| \geq q + 3$, we conclude that 3 divides q . Write $q = 3q_0$. Since $|s| \leq 1$, $|p| \leq 2$ and $|v| \geq q + 3$, we see that $|f(u)| \geq q$. Thus $f(u)$ has a prefix of length $q = 3q_0$ which is in $\ker \psi$. As the frequency matrix of f is invertible modulo 4, the prefix of u of length q_0 is in $\ker \psi$. We see that

$$\frac{|v|}{q} \leq \frac{3|u| + 3}{3q_0} = \frac{|u|}{q_0} + \frac{1}{q_0}.$$

Lemma 2. *Let s be a non-negative integer. If factor v of w has kernel period q , where $q \leq 1966(3^s)$, then*

$$\frac{|v|}{q} < \frac{35}{34} + \frac{3}{1966} \sum_{j=0}^{s-1} 3^{-j}.$$

Proof: If $s = 0$, this is implied by R2. Suppose $t > 0$ and the result holds for $s < t$. Suppose that $1966(3^{t-1}) < q \leq 1966(3^t)$ and there is a factor v of w such that v has kernel period q . Suppose that $|v|/q \geq 35/34$. Without loss of generality, suppose that no extension of v has period q . We have seen that there is a factor u of w with kernel period $q_0 = q/3$, $1966(3^{t-2}) < q_0 \leq 1966(3^{t-1})$ such that

$$\begin{aligned} |v|/q &\leq |u|/q_0 + 1/q_0 \\ &< \left(\frac{35}{34} + \frac{3}{1966} \sum_{j=0}^{t-2} 3^{-j} \right) + \frac{1}{q_0} \text{ (by the induction hypothesis)} \\ &< \frac{35}{34} + \frac{3}{1966} \sum_{j=0}^{t-2} 3^{-j} + \frac{1}{1966(3^{t-2})} \\ &= \frac{35}{34} + \frac{3}{1966} \sum_{j=0}^{t-2} 3^{-j} + \frac{3}{1966(3^{t-1})} \\ &= \frac{35}{34} + \frac{3}{1966} \sum_{j=0}^{t-1} 3^{-j}. \square \end{aligned}$$

Theorem 3. *Word w contains no factor v with kernel period q such that*

$$|v|/q \geq 35/34 + 9/2(1966).$$

Proof: Suppose that factor v of w has kernel period q such that (1) holds. By Remark 1, we have $q \geq 1966$. By the previous lemma, for some non-negative s ,

$$|v|/q < \frac{35}{34} + \frac{3}{1966} \sum_{j=0}^{s-1} 3^{-j} < \frac{35}{34} + \frac{3}{1966} \sum_{j=0}^{\infty} 3^{-j} = \frac{35}{34} + \frac{9}{2(1966)}. \square$$

Corollary 4. *Dejean's conjecture holds for $n = 30, 31, 32$.*

Appendix: Computer search

Suppose that some factor v of w has kernel period $q \leq 1966$ and either $31(|v|+1) \geq 32q-3$ or $|v|/q \geq 35/34 + 9/2(1967)$. Without loss of generality, taking such a v as short as possible, we may assume that

$$|v| \leq \left\lceil \frac{32(1966) - 3}{31} - 1 \right\rceil = 2029.$$

$$(\text{We also have } \left\lceil (1966) \left(\frac{35}{34} + \frac{9}{2(1967)} \right) \right\rceil = 2029.)$$

If $|v| > 3$, v is a factor of $f(u)$ for some factor u of w with $|u| \leq (|v| + 4)/3$. For a non-negative integer r , let $g(r) = \lfloor (r + 4)/3 \rfloor$. Since $g^7(2029) = 2 < 3$, (here the exponent denotes iterated function composition) word v must be a factor of $f^7(u)$ for some factor u of w , $|u| = 2$.

The word $u_0 = 23141121142$ contains all 8 factors of w which have length 2. To establish R1 and R2, one thus checks that they hold for the single word $f^7(u_0)$ (which is of length 24,057). In fact, we performed this computer check on the word $f^7(u_1)$, where $u_1 = 11421231211231411$ contains all 13 factors of w which have length 3.

References

- [1] F. J. Brandenburg, Uniformly growing k -th powerfree homomorphisms, *Theoret. Comput. Sci.* **23** (1983), 69–82.

- [2] J. Brinkhuis, Non-repetitive sequences on three symbols, *Quart. J. Math. Oxford* (2) **34** (1983), 145–149.
- [3] A. Carpi, On Dejean’s conjecture over large alphabets, *Theor. Comput. Sci.* **385** (2007), 137–151.
- [4] Françoise Dejean, Sur un théorème de Thue, *J. Combin. Theory Ser. A* **13** (1972), 90–99.
- [5] L. Ilie, P. Ochem, J. Shallit, A generalization of repetition threshold, *Theoret. Comput. Sci.* **345** (2005), 359–369.
- [6] D. Krieger, On critical exponents in fixed points of non-erasing morphisms, *Theoret. Comput. Sci.* **376** (2007), 70–88.
- [7] M. Lothaire, *Combinatorics on Words*, Encyclopedia of Mathematics and its Applications **17**, Addison-Wesley, Reading (1983).
- [8] F. Mignosi, G. Pirillo, Repetitions in the Fibonacci infinite word. *ITA* **26** (1992) 199–204.
- [9] M. Mohammad-Noori, J. D. Currie, Dejean’s conjecture and Sturmian words, *Euro. J. Combin.* **28** (2007), 876–890.
- [10] Jean Moulin-Ollagnier, Proof of Dejean’s conjecture for alphabets with 5, 6, 7, 8, 9, 10 and 11 letters, *Theoret. Comput. Sci.* **95** (1992), 187–205.
- [11] Jean-Jacques Pansiot, A propos d’une conjecture de F. Dejean sur les répétitions dans les mots, *Discrete Appl. Math.* **7** (1984) 297–311.
- [12] A. Thue, Über unendliche Zeichenreihen, *Norske Vid. Selsk. Skr. I. Mat. Nat. Kl. Christiana* No. 7 (1906).
- [13] A. Thue, Über unendliche Zeichenreihen, *Norske Vid. Selsk. Skr. I. Mat. Nat. Kl. Christiana* (1912), 1–67.